

MSL1 is a mechanosensitive ion channel that dissipates mitochondrial membrane potential and maintains redox homeostasis in mitochondria during abiotic stress

Chun Pong Lee¹, Grigory Maksaev², Gregory S. Jensen², Monika W. Murcha¹, Margaret E. Wilson², Mark Fricker³, Ruediger Hell⁴, Elizabeth S. Haswell², A. Harvey Millar¹ and Lee J. Sweetlove^{3,*}

¹ARC Centre of Excellence in Plant Energy Biology, University of Western Australia, Bayliss Building M316, 35 Stirling Highway, Crawley 6009, Western Australia, Australia,

²Department of Biology, Washington University in Saint Louis, One Brookings Drive, Mailcode 1137, Saint Louis, MO 63130, USA,

³Department of Plant Sciences, University of Oxford, South Parks Road, Oxford OX1 3RB, UK, and

⁴Department of Plant Molecular Biology, Centre for Organismal Studies, University of Heidelberg, Im Neuenheimer Feld 360, D-69120 Heidelberg, Germany

Received 24 May 2016; revised 4 August 2016; accepted 5 August 2016; published online 3 November 2016.

*For correspondence (e-mail lee.sweetlove@plants.ox.ac.uk).

SUMMARY

Mitochondria must maintain tight control over the electrochemical gradient across their inner membrane to allow ATP synthesis while maintaining a redox-balanced electron transport chain and avoiding excessive reactive oxygen species production. However, there is a scarcity of knowledge about the ion transporters in the inner mitochondrial membrane that contribute to control of membrane potential. We show that loss of MSL1, a member of a family of mechanosensitive ion channels related to the bacterial channel MscS, leads to increased membrane potential of Arabidopsis mitochondria under specific bioenergetic states. We demonstrate that MSL1 localises to the inner mitochondrial membrane. When expressed in *Escherichia coli*, MSL1 forms a stretch-activated ion channel with a slight preference for anions and provides protection against hypo-osmotic shock. In contrast, loss of MSL1 in Arabidopsis did not prevent swelling of isolated mitochondria in hypo-osmotic conditions. Instead, our data suggest that ion transport by MSL1 leads to dissipation of mitochondrial membrane potential when it becomes too high. The importance of MSL1 function was demonstrated by the observation of a higher oxidation state of the mitochondrial glutathione pool in *msl1-1* mutants under moderate heat- and heavy-metal-stress. Furthermore, we show that MSL1 function is not directly implicated in mitochondrial membrane potential pulsing, but is complementary and appears to be important under similar conditions.

Keywords: mechanosensitive ion channel, oxidative stress, mitochondria, membrane potential, pulsing, Arabidopsis.

INTRODUCTION

The bioenergetic function of mitochondria is crucially dependent on the impermeable nature of their inner membrane to protons such that an electrochemical gradient can be established by the proton-pumping activity of the respiratory chain. This provides the proton motive force (Δp) that powers the F_1F_0 ATP synthase. In plant mitochondria, the presence of an active K^+/H^+ exchange across the inner membrane (Hensley and Hanson, 1975; Zotova *et al.*, 2010) means that ΔpH is a relatively minor component of Δp in plant mitochondria (Mitchell and Moyle, 1969; Nicholls, 1974; Santo-Domingo and Demarex, 2012). In addition to

K^+/H^+ exchange, it has long been apparent that plant mitochondrial membranes are permeable to other cations and anions (Moore and Wilson, 1977; Huber and Moreland, 1979; Jung and Brierley, 1979). However, the transporter proteins responsible for these ion fluxes are largely unknown, and the purpose of these other ion-transport capabilities has not been established. One possibility is that ion transporters lead to a net transfer of charge from the intermembrane space to the matrix that serves to dissipate mitochondrial membrane potential ($\Delta\Psi_M$) and results in the endogenous inner membrane 'leak' observed in

energised mitochondria. This is apparent as continued electron transport (oxygen consumption) in the absence of dissipation of ΔpH by F_1F_0 ATP synthase, and can be stimulated by metals such as cadmium (Kessler and Brand, 1994a,b,c). While membrane leak is disadvantageous in terms of ATP production, it would be beneficial in situations where there is an imbalance between the rate of electron transport/proton pumping and F_1F_0 ATP synthase activity, leading to a very high electrochemical potential across the inner membrane. Such an imbalance leads to increased production of damaging reactive oxygen species (ROS) because the large electrochemical gradient makes it harder for the respiratory chain complexes to pump protons against that gradient. Because redox cycling of complexes I, III and IV is coupled to proton pumping, this has the effect of reducing the rate of electron transport in the respiratory chain. As a result, the respiratory complexes become highly reduced, which increases the propensity for ROS production by transfer of single electrons to molecular oxygen (Møller, 2001). Ion transport events that reduce the electrochemical gradient would therefore help prevent ROS production by easing the restriction on respiratory chain electron transport.

One established mechanism in which ion transport process functions in this way is mediated by uncoupling protein (UCP; Vercesi *et al.*, 2006) that allows movement of protons from the intermembrane space to the matrix, most likely via an indirect mechanism involving transport of anionic fatty acids (Jezek *et al.*, 1997). UCP is activated under conditions of bioenergetic imbalance (Smith *et al.*, 2004) and has been demonstrated to reduce the production of mitochondrial ROS and to protect against oxidative stress (Barreto *et al.*, 2014). Although activation of UCP is rapid, inactivation probably involves the turnover of the protein (Azzu *et al.*, 2010) and thus UCP can be considered a medium timescale response to persistent mitochondrial imbalance caused by environmental stress. However, to maintain homeostasis there must also be mechanisms that can operate on a short timescale. Rapid, short-term dissipation of $\Delta\Psi_M$ has been observed in plant mitochondria with transients (pulses) lasting less than 1 min (Schwarzländer *et al.*, 2012). This appears to involve selective cation transport, although the transporter or channel responsible has not been identified. Although the first molecular details of regulated cation transport in plant mitochondria are beginning to emerge with the identification of proteins involved in mitochondrial Ca^{2+} transport (Wagner *et al.*, 2015; 2016), it is not yet clear if these proteins are involved in bioenergetic balancing. More work needs to be done to fully understand the mechanisms of ion transport in plant mitochondria, and the biochemical and physiological roles of the transport events.

To this end, we searched for putative mitochondrial transporter proteins in Arabidopsis based on sequence

analysis (identification of a mitochondrial-targeting peptide and presence of extensive hydrophobic domains of sufficient length to be membrane spanning). One of the proteins we identified is a member of the MscS-Like (MSL) family of mechanosensitive ion channels (At4g00290; MSL1; Hamilton *et al.*, 2015a). This 10-member gene family was initially identified in Arabidopsis based on homology with the bacterial mechanosensitive ion channel of small conductance MscS (Pivetti *et al.*, 2003). The MSL proteins have different subcellular locations, and have been identified or predicted to localise to the plasma membrane, endoplasmic reticulum and plastid, with MSL1 being the only member with a putative mitochondrial localisation (Hamilton *et al.*, 2015a). Like their bacterial homologs, the MSL proteins characterised to date can be stretch-activated ion channels and show a slight preference for anions (Maksaev and Haswell, 2012; Hamilton *et al.*, 2015b). In bacteria, stretch-activation of MscS promotes survival of hypo-osmotic shock: opening of the channel following cell swelling leads to the release of ions, reducing the cell's osmotic potential (Levina *et al.*, 1999). A similar hypo-osmotic shock protection role within plastids and during pollen germination has been demonstrated for specific Arabidopsis MSL family members (Veley *et al.*, 2012; Hamilton *et al.*, 2015b). Nevertheless, MSL proteins also appear to have other functions in plants, including control of plastid division (Wilson *et al.*, 2011) and activation of programmed cell death by the plasma membrane-localised MSL10 (Veley *et al.*, 2014). In this paper, we present a characterisation of the localisation and electrophysiological properties of MSL1, and investigate the conditions under which it is important for bioenergetics homeostasis.

RESULTS

MSL1 is an integral membrane protein in the mitochondrion

Among the 10 MSL proteins that harbor a conserved MscS domain in Arabidopsis, only MSL1 has a consensus mitochondrial class II (-3 R) cleavage site at Phe-79 (RAF/SS; Huang *et al.*, 2009). It is also the MSL family member that has been identified by previous proteomic analyses in either mitochondria- or plastid-enriched fractions (Froehlich *et al.*, 2003; Zybailov *et al.*, 2008; Ferro *et al.*, 2010; Klodmann *et al.*, 2011; Nikolovski *et al.*, 2012), raising the possibility that MSL1 may be dual-targeted. To clarify the subcellular localisation of MSL1, a C-terminal MSL1-GFP fusion was expressed under the control of the 35S promoter in transgenic Arabidopsis plants. Confocal laser-scanning microscope (CLSM) imaging of leaf epidermal cells in this transgenic line revealed that MSL1-GFP localised to subcellular structures resembling mitochondria (Figure 1a). MSL1-GFP co-localised with the mitochondrial marker MitoTracker Red in root hair cells (Figure 1b),

confirming a mitochondrial localisation. To further verify MSL1 localisation, an *in vitro* import assay was carried out by incubating radiolabeled precursor MSL1 with isolated mitochondria under conditions that support import (Figure 1c). MSL1 was imported into a proteinase K-resistant location inside mitochondria. Import required a negative mitochondrial membrane potential, as evidenced by the lack of proteinase K-protected mature MSL1 in the presence of valinomycin (Figure 1c). The presence of a processing intermediate and a mature protein indicate that MSL1 is most likely imported via a two-step process common to the import of inner membrane proteins that contain a cleavable pre-sequence (Murcha *et al.*, 2004). In contrast, MSL1 is not imported into chloroplasts *in vitro* (Figure S1), providing additional evidence that MSL1 is exclusively localised to mitochondria.

MSL1 is reported to be uniformly expressed across tissue types and developmental stages, as well as under most biotic and abiotic treatments (Genevestigator and

AtGenExpress Visualisation Tool analysis). To analyze the physiological and functional relevance of MSL1 in plants, we isolated a homozygous T-DNA insertion line (*msl1-1*) that carries a T-DNA insertion in the first exon leading to an absence of *MSL1* transcript (Figure S2a,b). A homozygous *msl1-1* line complemented with a construct containing the genomic version of *MSL1* controlled by its native promoter was also generated (*msl1-1/gMSL1*; Figure S2b, c). Both transgenic lines show similar growth and morphology to wild-type (WT) plants under standard growth conditions on agar plates and on soil (Figure S2d). Using these genetic resources we confirmed that a peptide antibody raised against the C-terminal end of MSL1 was specific for this protein (Figure S2c). We then further used this antibody to assess the precise intra-mitochondrial localisation and orientation of MSL1. Based on the consensus TM prediction algorithms in ARAMEMNON (Schwacke *et al.*, 2003), MSL1 is probably comprised of five transmembrane helices. When isolated mitochondrial membranes were

Figure 1. The subcellular localisation of MSL1 in Arabidopsis.


(a) Confocal laser-scanning microscope (CLSM) image of mesophyll cells from transgenic plants stably over-expressing MSL1-GFP. GFP signal is pseudocolored green and chlorophyll fluorescence is pseudocolored red. Scale bar: 20 μ m.

(b) CLSM image of a root hair from a plant stably over-expressing MSL1-GFP after incubation with MitoTracker Red. GFP signal is pseudocolored green and MitoTracker red is pseudocolored red. Scale bar: 2 μ m.

(c) *In vitro* import of radiolabeled MSL1 into isolated mitochondria. Import reaction was carried out under conditions that support protein uptake into mitochondria. p, precursor protein; i, processed intermediate; m, processed mature protein.

(d) Immunoblotting of isolated mitochondria and mitochondrial peripheral and integral membrane fractions following sodium carbonate treatment. Proteins were separated on a blue-native PAGE and immunoblots were probed using a peptide antibody raised against MSL1. Roman numerals correspond to the locations of respiratory complexes and ATP synthase complex on a typical 1D-blue-native PAGE of Arabidopsis mitochondria.

(e) Sub-mitochondrial location of MSL1. Isolated mitochondria were treated with (+) or without (-) osmotic shock to produce mitoplasts, proteinase K to digest exposed peptides and triton (TX-100) to solubilise all membranes. Immunoblotting was then carried out using antibodies raised against MSL1, stomatin 2 (SLP2, inner membrane), voltage-dependent anion channel (VDAC, outer membrane) and cytochrome c (cyt c, intermembrane space).



fractionated into carbonate-soluble (peripheral) and -insoluble (integral) membrane proteins and analyzed by Blue-Native PAGE and immunoblotting, MSL1 could only be found in the integral membrane fraction and migrated as a large ~350-kDa oligomeric complex (after correction by a factor of 0.7 for digitonin and Coomassie binding; Wittig *et al.*, 2010; Figure 1d). The immunoreactive protein band on Blue-Native PAGE was confirmed to be MSL1 by analysis of trypsin-digest-peptides using mass spectrometry (Data S1). Furthermore, no trypsin-cleavable peptide fragments preceding Ser-83 (i.e. the mitochondrial targeting sequence) were detected (Data S1), suggesting that MSL1 is imported into the mitochondrial matrix where its pre-sequence is cleaved before it is inserted, most likely, into the inner membrane. To test this, we assessed the accessibility of MSL1 to degradation by proteinase K in inner membrane vesicles lacking the outer membrane (mitoplasts) by immunoblotting of MSL1 and marker proteins. Successful preparation of outer membrane-free mitoplasts was confirmed by the absence of the voltage-dependent anion channel (VDAC, outer membrane) and the sensitivity of cytochrome c (intermembrane space) to degradation by proteinase K (Figure 1e). Both an MSL1-containing complex and the inner membrane protein SLP2 (Gehl *et al.*, 2014) were detected in the mitoplast fraction treated with proteinase K (Figure 1e). The detection of an MSL1 band by anti-MSL1 antibodies, which recognise a short hydrophilic peptide sequence near the C-terminal end of MSL1 (see Experimental procedures), indicates that this region of the protein is in the matrix and is not accessible to proteinase K. In summary, MSL1 is an integral inner membrane protein of mitochondria with the C-terminus orientated towards the matrix.

MSL1 activity is a stretch-activated ion channel

To establish if MSL1 forms a mechanosensitive ion channel, we used single-channel patch-clamp electrophysiology. An N-terminal 6xHis-*AtMSL1* construct was expressed in an *Escherichia coli* strain that lacks all seven known mechanosensitive ion channels (MJF641, $\Delta 7$; Edwards *et al.*, 2012) to provide an electrophysiologically silent background. When expressed in *E. coli*, the 6xHis-tagged mature MSL1 (lacking its predicted mitochondrial targeting peptide) can be found in the triton-soluble fraction (Figure S3a) and forms a large complex with a similar oligomeric size to the mitochondrial MSL1 (Figures 1d and S3b). Giant *E. coli* spheroplasts expressing mature MSL1 produced ion channel activity in response to increased lateral membrane tension, applied as suction – or negative pressure – through the patch pipette (Figure 2a).

The current/voltage plot for MSL1 gated by tension between +40 and –180 mV (Figure 2b) reveals that at negative membrane potentials, MSL1 has a single channel conductance of 1.2 nS. This is the same single channel

conductance observed for *E. coli* MscS in the same system (Kung *et al.*, 2010; Booth *et al.*, 2011). However, MSL1 activity in *E. coli* spheroplasts was less stable than *Ec* MscS, alternating more frequently between conducting and non-conducting states. MSL1 is likely to be oriented in the *E. coli* plasma membrane the same way as *Ec* MscS, with the C-terminus facing the cytoplasm (Miller *et al.*, 2003). MSL1 is similarly oriented in the inner membrane of mitochondria, with the C-terminus in the matrix (Figure 1). Both the *E. coli* plasma membrane and the mitochondrial inner membrane have strongly negative membrane potentials, indicating that MSL1 is likely to have high conductance in mitochondria once it is gated by membrane tension. Even in the absence of applied tension, we routinely observed noisy activity that may represent MSL1 channel opening at strongly negative membrane potentials (–160 mV or higher, double asterisk in Figure 2c). However, the membranes of *E. coli* giant spheroplasts deteriorate under such high membrane potentials, and it is therefore formally possible that this activity represents current fluctuation associated with imminent patch rupture (Figure 2c, arrow). We occasionally (in about 20% of patches) observed more easily identified opening and closing events without added tension at moderate membrane potentials (–60 to –100 mV; Figure 2c, single asterisk), suggesting that MSL1 may be voltage-gated, although it remains possible that this activity may be a response to transient fluctuations in patch tension. At positive membrane potentials, which are not physiologically relevant for *E. coli* or for the inner mitochondrial membrane, MSL1 channel activity was reduced and unstable (Figure S4).

MSL1 was found to have a very mild preference for anions (Figure 2d), with a reversal potential (E_{rev}) of 10.2 mV and a preference for anions over cations ($P_{Cl}:P_K$) by a factor of approximately 2.2. These characteristics closely resemble those reported previously for *Ec*MscS under the same experimental conditions in giant spheroplasts [$E_{rev} = 9.5$ mV (Edwards *et al.*, 2008), $E_{rev} = 12$ mV (Li *et al.*, 2002)] and when *Ec*MscS was reconstituted into the liposomes [$P_{Cl}:P_K = 1.5$ (Sukharev, 2002); $P_{Cl}:P_K = 2.7$ (Cox *et al.*, 2013)]. These results indicate that MSL1 is essentially non-selective and when open allows the passage of small anions and cations with similar efficiency.

MSL1 can provide modest protection against hypo-osmotic shock in *Escherichia coli* but not in plant mitochondria

In bacteria, MscS protects against hypo-osmotic shock; the increased membrane tension due to cell swelling causes stretch-activated opening of the channel and release of ions, reducing the osmotic potential of the cell (Naismith and Booth, 2012). To determine if MSL1 was similarly capable of protecting bacterial cells from hypo-osmotic shock, MSL was expressed in the *E. coli* strain MJF465 that lacks




Figure 2. Channel function of MSL1 when expressed in *Escherichia coli*. (a) Representative trace of an inside-out patch from a giant $\Delta 7$ (MJF641) *E. coli* spheroplast expressing mature MSL1 at a transmembrane potential of -20 mV. Channels were opened by administering negative pressure (suction) through the patch pipette. O_1 – O_3 indicate successive individual channel openings. (b) The current–voltage curve for MSL1 at negative membrane potentials. The single channel conductance of MSL1, calculated as the slope of this I/V curve, is 1.2 nS. $N = 17$ patches. (c) The same patch shown in (a) subjected to extreme transmembrane potentials without added membrane tension. The asterisk indicates occasional channel openings and closures. The arrow indicates patch rupture. (d) Current–voltage properties of the MSL1 channel under symmetric (filled circles, data are the same as in b) and asymmetric (open circles) conditions. Arrow marks the reversal potential of MSL1 in asymmetric buffer.

three key mechanosensitive ion channels, MscS, MscL and MscK (Levina *et al.*, 1999). As expected, MJF465 cells do not survive severe hypo-osmotic shock, but this sensitivity could be fully rescued by expressing MscS from an IPTG-inducible plasmid (Figure 3a). Inducible expression of MSL1 resulted in partial rescue (40% of WT survival) of the MJF465 osmotic shock sensitivity. Similar results were previously obtained for MSL3, a plastid-localised MscS homolog also from *Arabidopsis* (Haswell and Meyerowitz, 2006). To establish if MSL1 fulfils a similar osmotic shock protection role in plant mitochondria, we examined the hypo-osmotic swelling behavior of mitochondria isolated from WT and *msl1-1* seedlings. After labeling with the membrane-potential-independent dye MitoTracker Green (MTG), mitochondria were subjected to hypo-osmotic shock and were visualised under CLSM (Figure 3b). Mitochondrial volumes were calculated from Z-stacks (Figure 3c), and it is apparent that as the concentration of osmoticum decreases, the mitochondria swell. However, the degree of swelling was not significantly different between WT and *msl1-1*. When osmoticum was completely omitted, the majority of mitochondria appeared ruptured, but the number of intact mitochondria that remained was the same in *msl1-1* and WT mitochondria. These data demonstrate that MSL1 does not provide substantial protection against rupture from hypo-osmotic shock in isolated mitochondria.

Loss of MSL1 leads to increased membrane potential in isolated mitochondria when F_1F_0 ATP synthase is inactive

Given that MSL1 is essentially non-selective with respect to ion species (Figure 2d) and that these channels are not known to exclude H^+ or K^+ (Maksaev and Haswell, 2012; Naismith and Booth, 2012; Cox *et al.*, 2013; Hamilton *et al.*, 2015b), opening of MSL1 would be expected to lead to partial dissipation of the mitochondrial membrane potential ($\Delta\Psi_M$). We therefore sought to determine the effect of MSL1 on $\Delta\Psi_M$ of isolated mitochondria and to use this readout to explore the conditions under which MSL1 is active. Isolated mitochondria from WT and *msl1-1* and the complemented line, *msl1-1/gMSL1*, were energised with respiratory substrates and labeled with MTG and the cationic lipophilic fluorescent molecule tetramethyl rhodamine methyl ester (TMRM). The use of these two fluorescent probes provides a ratiometric estimation of membrane potential given that accumulation of TMRM in mitochondria is $\Delta\Psi_M$ -dependent, while MTG accumulates in a $\Delta\Psi_M$ -independent manner, providing a reference control. TMRM and MTG fluorescence were quantified by CLSM imaging (Figure 4a). When mitochondria were energised with succinate, a significantly higher proportion of *msl1-1* mitochondria had a higher TMRM:MTG fluorescence ratio than WT and the complemented line (Figure 4b), demonstrating that




Figure 3. Function of MSL1 during osmotic shock.

(a) Osmotic shock survival of *Escherichia coli* expressing AtMSL1. Osmotic shock assay with WT (Frag-1) or *mscL*-*mscS*-*mscK*- (MJF465) *E. coli* strains harboring the indicated constructs. Survival of cells subjected to a 500 mM hypo-osmotic downshock is presented as a percentage of the survival of unshocked cells. Data shown are the average of three different experiments and error bars indicate standard deviation. Asterisk indicates significant difference ($P < 0.01$) by Student's *t*-test.

(b) Confocal laser-scanning microscope (CLSM) image showing the maximum intensity projection of mitochondria isolated from Col-0 and *msl1-1* following osmotic treatments. Isolated mitochondria were labeled with membrane potential-independent fluorescent dye MitoTracker Green, and were then subjected to osmotic shock. Labeled mitochondria were visualised by confocal microscopy with at least 10 focal steps (z-stacks) of 0.2 μm . Scale bar: 1 μm .

(c) Volume of isolated mitochondria from (b). The volume of each mitochondrion was estimated on the basis of the maximum intensity projection-derived radius, with the assumption that every mitochondria visualised after all *in vitro* treatments remain perfectly spherical. Data shown are the average of three different experiments, with at least 200 mitochondria analyzed per experiment, and error bars denote standard deviation. Asterisk indicates significant difference ($P < 0.05$) between sucrose treatments but not between genotypes according to the two-way ANOVA analysis.


loss of MSL1 leads to higher $\Delta\Psi_M$ in the presence of respiratory substrate but no ADP. Similar results were obtained when malate and pyruvate were used as respiratory substrates (Figure 4c), suggesting that under state II conditions when the F_1F_0 ATP synthase is inactive due to lack of ADP and the $\Delta\Psi_M$ is thus high, the MSL1 channel is active in WT and complemented lines. This would serve to limit the maximum membrane potential that can be reached.

To establish whether the bioenergetic state of the mitochondrion influences MSL1 function, we repeated the analysis with mitochondria in state III (supplied with succinate and ADP). As expected, the activation of the F_1F_0 ATP synthase caused a decrease in $\Delta\Psi_M$ measured as a decrease in the TMRM:MTG fluorescence ratio. Additionally, it was apparent that there was no longer a difference in the TMRM:MTG ratio between the genotypes (Figure 4d). To

Figure 4. Effect of the loss of MSL1 on membrane potential of isolated mitochondria in different respiratory states.

(a) Representative confocal image of isolated mitochondria from Col-0, *msl1* and complemented line (*msl1/gMSL1*) incubated with minimal respiration buffer containing succinate to induce state II respiration and loaded with fluorophores TMRM and MTG. TMRM is false-colored red and MTG is false-colored green. Scale bar: 1 μm .

(b–e) Frequency distribution of $\Delta\Psi_M$ (expressed as the ratio between log-transformed intensity of TMRM and MTG) of isolated mitochondria in the presence of: (b) succinate (state II respiration); (c) malate and pyruvate (state II respiration); (d) succinate and 0.5 mM ADP (state III respiration); (e) succinate, 0.5 mM ADP and 1 μM oligomycin. Data shown are the average of at least four different experiments, with at least 500 mitochondria analyzed per experiment, and error bars denote standard error of the mean. Asterisk on each panel label indicates significant difference ($P < 0.01$ according to ANOVA and Tukey's *post hoc* analysis) between Col-0 versus *msl1-1* and *msl1-1* versus *msl1-1/gMSL1* (complemented line) in the mean of TMRM:MTG fluorescence ratio from at least four independent experiments.



confirm that the genotype response under state III conditions was due to ATP synthase activity, oligomycin was added to inhibit ATP synthase and this restored the difference in TMRM:MTG ratio between genotypes (Figure 4e). Hence, loss of MSL1 leads to an increased $\Delta\Psi_M$ compared with WT, but only under state II conditions when F_1F_0 ATP synthase is inactive. Given that opening of a non-selective ion channel such as MSL1 would dissipate $\Delta\Psi_M$, this is likely a direct consequence of the absence of MSL1, although an indirect or secondary effect cannot be ruled out.

MSL1 is not responsible for $\Delta\Psi_M$ pulsing, but loss of MSL1 creates conditions that increase pulsing

Some of the frequency distribution histograms of the TMRM:MTG fluorescence ratio in individual isolated

mitochondria taken at single time-points (Figure 4) appear bimodal. One possible cause of bimodal populations of $\Delta\Psi_M$ is membrane potential pulsing, which is prevalent in isolated mitochondria in state II (Schwarzlander *et al.*, 2012). These $\Delta\Psi_M$ pulses are frequent in state II (on a time-scale of seconds to minutes), and rather irregular in frequency and amplitude (Schwarzlander *et al.*, 2012). Hence, in a snapshot of a population of mitochondria (as shown in Figure 4), some mitochondria will be between pulses (higher $\Delta\Psi_M$) and some will be in a pulse (lower $\Delta\Psi_M$), leading to two peaks in the TMRM:MTG frequency distribution histograms. Moreover, the state II bioenergetic condition that promotes pulsing is the same condition under which loss of MSL1 affects $\Delta\Psi_M$, raising the possibility that transient opening of MSL1 could be responsible for pulsing. To investigate this further, we quantified $\Delta\Psi_M$ (from

TMRM fluorescence) over a time-course in multiple individual isolated state II mitochondria from WT and *msl1-1* and the complemented line (Figure 5). As expected, state II WT mitochondria showed frequent $\Delta\Psi_M$ pulses (Figure 5a). However, frequent $\Delta\Psi_M$ pulsing was also apparent in state II *msl1-1* mitochondria, meaning that MSL1 is not the channel responsible for pulsing. Frequency distribution histograms of the complete set of these time-course data were plotted for populations of mitochondria from WT, the *msl1-1* knockout and the complemented line (Figure 5b) and, as in the single time-point snapshots of Figure 4, a bimodal distribution was apparent that could be fitted with a two-component Gaussian mixture model. The same analysis was done for mitochondria from the three genotypes under state III (in the presence of ADP). As expected this dramatically reduced the extent of pulsing in all genotypes (Figure 5c) and decreased $\Delta\Psi_M$ (Figure 5d). We previously established that the best statistical measure of the amount of pulsing is given by the coefficient of variation (CV) of $\Delta\Psi_M$ over time for each mitochondrion (Schwarzl ander *et al.*, 2012). The mean CV from multiple mitochondria revealed that the amount of pulsing was significantly greater in the *msl1-1* mutant (Figure 5e). Additionally, the proportion of the population of mitochondria that was pulsing (given by the area under the left-hand peak in the TMRM:MTG frequency distribution histogram with decreased $\Delta\Psi_M$) was also significantly greater in the *msl1-1* mutant (Figure 5g), suggesting a greater frequency or duration of pulsing. Finally, to provide a robust measure of the resting $\Delta\Psi_M$ (i.e. between pulses) and to confirm the apparent increase in $\Delta\Psi_M$ in the *msl1-1* line in state II mitochondria shown in Figure 4, we determined the mean TMRM:MTG value of the right-hand peak of the frequency distribution corresponding to the higher $\Delta\Psi_M$ that occurs between pulses. This confirmed a significant increase in $\Delta\Psi_M$ in the *msl1-1* mutant in comparison to the WT and complemented line (Figure 5g), consistent with the data from single time-points (Figure 4). In summary, we infer that MSL1 is not directly responsible for $\Delta\Psi_M$ pulsing, but that loss of MSL1 does increase pulsing frequency and amplitude. The most likely role of MSL1 is thus to help to limit build-up of excessive membrane potential during temporary bioenergetic imbalance. Likewise, loss of MSL1 may exacerbate the bioenergetic


conditions that cause pulsing (high membrane potential and ROS production).

The effect of MSL1 on $\Delta\Psi_M$ in isolated mitochondria is moderated by ROS scavengers and antioxidants

When mitochondria are energised by respiratory substrate but ATP synthase is not active (state II), the respiratory chain becomes highly-reduced and the rate of mitochondrial ROS production increases (M oller, 2001). This phenomenon and the changes in pulsing (Figure 5) raise the possibility that MSL1 channel function is influenced by the redox status of mitochondria. To test this, we incubated state II mitochondria with a range of different antioxidants, free radical scavengers and lipid peroxidation inhibitors (Figure 6). If MSL1 is directly activated by the ROS produced in state II mitochondria, then we would expect the addition of ROS scavengers to increase WT membrane potential to mutant values, i.e. the prevention of ROS activation of MSL1 under state II conditions would mean that it no longer serves to dissipate $\Delta\Psi_M$. Thus, by preventing activation of MSL1 by ROS, the WT could be expected to phenocopy the knockout mutant. This behavior is what was observed when the matrix-accumulating antioxidant MitoTEMPOL was added. However, a less clear picture emerged when other ROS-ameliorating agents were used. For example, addition of MitoQ, which specifically prevents inner membrane lipid peroxidation (Kelso *et al.*, 2001; Murphy and Smith, 2007), abolished the difference in $\Delta\Psi_M$ between *msl1-1* and WT, but not by increasing WT $\Delta\Psi_M$ to the level expected in the *msl1-1* mutant but by reducing $\Delta\Psi_M$ in the *msl1-1* mutant to WT levels. This was not the result of a non-specific uncoupling effect of MitoQ because the control analog decyl triphenylphosphonium (dTPP), which lacks antioxidant activity, did not alter $\Delta\Psi_M$ in the *msl1-1* mutant. The antioxidant, ascorbate, also had a similar effect to MitoQ, whereas the matrix-accumulating superoxide dismutase mimetic MnTMPyP had no effect. These data suggest that alterations in mitochondrial ROS can affect MSL1 activation state, but there is also evidence that loss of MSL1 alters the sensitivity of mitochondria to the antioxidants MitoQ and ascorbate. The interaction between MSL1, ROS and $\Delta\Psi_M$ is complex, and appears to depend on the local origin and type of ROS produced. At this stage we do not

Figure 5. Effect of MSL1 knockout on $\Delta\Psi_M$ pulsing of isolated mitochondria.

TMRM- and MTG-labeled mitochondria were energised either by succinate (state II respiration) (a, b) or succinate and ADP (state III respiration) (c, d), and the responses from these fluorophores in individual mitochondria over 2 min were monitored by confocal microscopy. MTG-normalised TMRM fluorescence intensity traces of five representative mitochondria (a, c) and ln (TMRM/MTG) frequency distribution of the population (b, d) from Col-0 (left), *msl1-1* (middle) and complemented line *msl1-1/gMSL1* (right) are shown. Gaussian fits to the frequency distributions are shown as colored lines (b, d), and where more than one Gaussian was fitted, the red line indicates the amalgamated fit and the green and blue lines the individual fits (b). The following statistics were extracted to quantify pulsing and membrane potential: (e) the coefficient of variation as a measure of pulsing amplitude; (f) the normalised TMRM fluorescence of state II mitochondria between pulses (the mean of the right-hand peak in the frequency distributions); (g) the proportion of pulsing (calculated as the normalised area under the curve of the left-hand peak in the frequency distribution plots). Each of these statistics was derived from at least 100 mitochondria in three independent experiments. Error bars indicate standard error and asterisks denote significant differences between the indicated pairs based on ANOVA and Tukey's *post hoc* analysis. * $P < 0.05$; ** $P < 0.01$; *** $P < 0.001$.






Figure 6. Effect of ROS scavengers and antioxidants on membrane potential of state II mitochondria isolated from WT, *msl1-1* and *msl1-1/gMSL1* plants. TMRM- and MTG-labeled mitochondria isolated from the three genotypes were energised with succinate (state II), and the effect of addition of the indicated ROS scavengers and antioxidants on TMRM and MTG fluorescence monitored by confocal microscopy. Each measurement consisted of at least 100 mitochondria and was repeated two-three times. For each measurement, a frequency distribution of TMRM:MTG was plotted and two Gaussian curves fitted to the bimodal distribution. The mean TMRM:MTG value of the right-hand of the two peaks was calculated. dTPP, decyl-triphenylphosphonium bromide; MnTMPyP manganese tetrakis(1-methyl-4-pyridyl)porphyrin pentachloride. Values are means \pm standard error, and asterisks denote significant differences between the indicated pairs based on ANOVA and Tukey's *post hoc* analysis. * $P < 0.05$; ** $P < 0.01$; *** $P < 0.001$.

have a clear understanding of the mechanism behind these interacting indirect effects.

MSL1 is crucial for regulating mitochondrial redox status during abiotic stress

So far we have demonstrated that the presence of MSL1 is important under bioenergetic conditions in which the respiratory chain is likely to become highly reduced and the rate of mitochondrial ROS production would increase. This bioenergetic state can often be triggered by abiotic stresses due to disruption of normal electron transport or general alterations in redox and energy balance (Møller, 2001). To establish whether MSL1 is important during abiotic stress, we monitored the *in vivo* mitochondrial redox status of WT and *msl1-1* roots expressing a mitochondria-targeted roGFP2 that reports on the redox poise of the mitochondrial glutathione pool (mito-roGFP2; Schwarzländer *et al.*, 2016) subjected to various pharmacological treatments and stresses (Figure 7). Figure 7(a) shows false-colored images of the ratio of mito-roGFP2 fluorescence collected with 405 and 488 nm excitation wavelengths. An increase in this ratio corresponds to greater oxidation of the roGFP2 probe and indicates greater oxidation state of the glutathione pool in the mitochondrial matrix (Schwarzländer *et al.*, 2016). As expected, incubation of roots with oligomycin to inhibit mitochondrial ATP synthase increased the oxidation of mito-roGFP2 in WT mitochondria (Figure 7a,b), because the reduced ATP synthase activity leads to over-reduction of respiratory chain complexes and subsequent ROS production in the

mitochondrial inner membrane. The increase in roGFP2 oxidation in response to oligomycin treatment was substantially greater in the *msl1-1* mutant than in WT, demonstrating that loss of MSL1 compromises the ability of mitochondria to maintain redox homeostasis *in vivo*. The involvement of mitochondrial bioenergetic imbalance was confirmed by the addition of a membrane uncoupler, carbonyl cyanide *m*-chlorophenyl hydrazone (CCCP), which restored mito-roGFP2 oxidation state to WT values (Figure 7b). The involvement of ROS was confirmed by the restoration of *msl1-1* mito-roGFP2 oxidation state to WT values by the addition of ascorbate (Figure 7b). Addition of antimycin A, which is known to raise mitochondrial membrane potential and generate mitochondrial ROS by inhibition of the Q cycle in complex III (Quinlan *et al.*, 2011), also caused an increase of mito-roGFP2 oxidation state in *msl1-1*, but rotenone, an inhibitor of complex I, had no effect (Figure 7b). In contrast, addition of propranolol, which prevents the uptake of inorganic anions as well as di- and tri-carboxylates by the plant inner mitochondrial membrane anion channel (PIMAC; Beavis, 1992; Laus *et al.*, 2008), reduced the roGFP2 oxidation state to a similar degree in WT and *msl1-1*.


We also examined the mito-roGFP2 oxidation state in roots exposed to a range of stress conditions, namely salt-, osmotic-, heavy-metal- and high-temperature-stress (Figure 7c). As found previously (Schwarzländer *et al.*, 2012), salt stress did not affect the oxidation of a mitochondria-targeted roGFP2 in WT roots, and there was no significant difference between WT and *msl1-1*. Similarly, there was no

Figure 7. Effect of absence of MSL1 on *in vivo* mitochondrial redox status during abiotic stress.

(a) Representative ratiometric pseudocolored images of mito-roGFP2 in Col-0 and *msl1-1* roots treated with oligomycin and Cd²⁺. roGFP2 fluorescence was measured for excitation at 405 and 488 nm, and the degree of oxidation is calculated based on the ratio between the intensities from the two excitations (405/408 ratio) and the calibration of the probe using 10 mM DTT and 10 mM H₂O₂. The color scale represents the redox state of roGFP2 ranging from fully reduced in indigo to fully oxidised in red. Scale bar: 50 µm.

(b) Quantitation of mito-roGFP2 oxidation state in Col-0 and *msl1-1* roots treated with respiratory inhibitors. Concentration of inhibitors used: 10 µM oligomycin A, 20 µM antimycin A, 20 µM rotenone, 1 mM sodium ascorbate, 1 mM propranolol and 25 µM carbonyl cyanide m-chlorophenyl hydrazone (CCCP).

(c) Quantitation of mito-roGFP2 oxidation state in Col-0 and *msl1-1* roots under abiotic stress conditions. All treatments were carried out in the dark for 1 h in ½MS medium (pH 5.7) unless otherwise stated. At least 10 seedlings in several independent experiments were examined, with error bars indicate standard error. 'a' and 'b' indicate significant difference ($P < 0.01$ based on ANOVA and Tukey's *post hoc* analysis), respectively, between Col-0 and *msl1-1* under the same treatment, and between control and treatment.



effect of absence of MSL1 on mito-roGFP2 following hyper-osmotic stress (150 and 300 mM mannitol treatments), with both a slight decrease in mito-roGFP2 oxidation in both WT and *msl1-1* roots. This corroborates the osmotic shock assays on isolated mitochondria (Figure 3). In contrast, high temperature (44°C for 10 min) caused a significant increase in mito-roGFP2 oxidation in WT roots (Figure 7c), and this was significantly exacerbated in *msl1-1* roots. At a more extreme temperature (48°C for 5 min), roGFP2 became more oxidised (approximately 60% oxidised), and the difference between *msl1-1* and WT disappeared (Figure 7c). There was also a significant increase in mito-roGFP2 oxidation in *msl1-1* compared with WT roots treated with 3 or 5 mM Cd²⁺. Collectively, these data demonstrate that loss of MSL1 exacerbates mitochondrial

glutathione pool oxidation under specific abiotic stresses, and suggest that MSL1 may provide a mechanism for maintenance of mitochondrial redox status under stress.

DISCUSSION

Our data demonstrate that MSL1 is a stretch-activated non-selective ion channel (Figure 2). Stretch activation provides a regulatory mechanism that allows the channel to respond to cell/organellar swelling due to hypo-osmotic stress. The resultant release of ions reduces osmotic potential and can protect against hypo-osmotic shock, which is thought to be an important function of MscS homologues in *E. coli* and in Arabidopsis plastids and pollen grains. And although *E. coli*-expressing MSL1 show enhanced tolerance of hypo-osmotic shock, we did not find

strong evidence that MSL1 fulfils this role in Arabidopsis mitochondria (Figure 3). In isolated mitochondria, reduction of the osmotic potential of the incubating medium had only a minor effect on WT mitochondrial volume, indicating a robust mechanism for volume homeostasis. The absence of MSL1 may have had no significant effect on this response because the osmotic shock we administered was too severe or too rapid for MSL1 to have an effect, or possibly due to redundant osmoregulatory mechanisms, as in *E. coli* (Levina *et al.*, 1999). In chloroplasts, osmotic pressure and ion homeostasis are controlled not only by MSL2 and MSL3 (Veley *et al.*, 2012), but also by a family of K⁺/H⁺ antiporters (KEA) that work cooperatively with other cation channels (Kunz *et al.*, 2014). A similar K⁺/H⁺ antiporter (KHE) also exists in plant mitochondria, although its molecular identity remains unknown to date (Hensley and Hanson, 1975; Dioloz and Moreau, 1985; Trono *et al.*, 2015). Nevertheless, it is also notable that treatment of WT and *msl1-1* roots with hyper-osmotic concentrations of mannitol did not cause a significant increase in oxidation of the mitochondrial glutathione pool (Figure 7c), suggesting that MSL1 is not a major part of the mechanism that maintains mitochondrial redox poise under this stress condition. Instead, our data point to MSL1 being important under conditions of bioenergetic imbalance characterised by high $\Delta\Psi_M$ and a highly reduced respiratory chain leading to ROS production (Figures 4 and 5). The increased impact of high temperature and Cd²⁺ treatments on the oxidation state of mitochondrially targeted roGFP2 in roots of *msl1-1* knockout mutants (Figure 7c) demonstrates that MSL1 is an important component of the molecular machinery that maintains mitochondrial redox status under these specific stresses. Both of these stresses are known to cause bioenergetic imbalances in mitochondria (Lin and Markhart, 1990; Kessler and Brand, 1994a,b,c; Atkin *et al.*, 2002) and also stimulate mitochondrial pulsing (Schwarzländer *et al.*, 2012). Indeed, in isolated mitochondria, MSL1 functions in the same bioenergetic conditions that lead to pronounced pulsing, whilst a lack of MSL1 enhances the extent of pulsing, suggesting that the two mechanisms are complementary. Given that opening of MSL1 would cause dissipation of $\Delta\Psi_M$ it is likely that, similar to pulsing, MSL1 acts as a release valve to avoid over-reduction of the respiratory chain and ROS production due to the build-up of a very high $\Delta\Psi_M$.

Clearly, an important feature of such an uncoupling ion channel would be that it selectively opens only when the mitochondria are in this unbalanced bioenergetic state. One possibility is that the channel is gated or modulated by high negative transmembrane voltage. Although we saw some evidence of this in patch-clamped *E. coli* spheroplasts expressing MSL1, the instability of spheroplast membranes at high negative voltages made it impossible to characterise this behavior in more detail. An alternative possibility

is that high $\Delta\Psi_M$ leads to changes in membrane tension and this leads to activation of MSL1. This could be mediated by damage to membrane lipids by ROS, leading to altered conformation and tension of the membrane. A final possibility is that MSL1 could be directly activated by ROS via oxidation of amino acid side groups in such a way that the membrane tension threshold for channel opening is altered. (Ridone *et al.*, 2015). We explored the effect of reducing ROS *in vivo* using mitochondrially targeted antioxidants (Figure 6). In one case (MitoTEMPOL), we found effects consistent with ROS activation of MSL1 in that MitoTEMPOL increased $\Delta\Psi_M$ in WT mitochondria to levels observed in the *msl1-1* mutant. Other antioxidants also abolished the differences in state II mitochondrial membrane potential between WT and the *msl1-1* mutant, but via unexplained indirect effects that operated in the *msl1-1* mutant background rather than in WT plants with functional MSL1. These differences could be related to the different mode of action of the antioxidant reagents used. MitoTEMPOL consists of an antioxidant piperidine nitroxide group conjugated to the lipophilic cation trimethylphosphonium that delivers the nitroxide group to the matrix face of the inner membrane in a membrane-potential-dependent fashion (Murphy and Smith, 2007). The nitroxide group can catalyse the dismutation of superoxide and detoxify Fe²⁺ ions (Samuni *et al.*, 1990; Bar-On *et al.*, 1999; Trnka *et al.*, 2009), thereby preventing Fenton chemistry that leads to hydroxyl radical formation. In contrast, MitoQ delivers the antioxidant ubiquinone into the hydrophobic core of the inner membrane preventing lipid peroxidation (Murphy and Smith, 2007), while ascorbate detoxifies hydrogen peroxide via matrix-localised ascorbate peroxidase activity. The contrasting effects of these antioxidants may suggest that a direct oxidation of the MSL1 protein by hydroxyl radicals may be a potential activating mechanism rather than alteration of membrane tension via lipid peroxidation, but more detailed studies would be required to establish this.

MSL1 now sits alongside a growing cast of players that act to uncouple plant mitochondria. These include mitochondrial membrane potential pulsing (Schwarzländer *et al.*, 2012), UCP (Vercesi *et al.*, 2006), and alternative respiratory pathways generated by electron flow via the alternative NAD(P)H dehydrogenases (Rasmusson *et al.*, 2004) and alternative oxidase (Moore *et al.*, 2013). Moreover, it is clear that MSL1, pulsing and UCP all respond in various ways to mitochondrial ROS and/or mitochondrial membrane lipid peroxidation. The effect of activation of these three ion transport processes is the same – to dissipate $\Delta\Psi_M$ – leading to the question of why more than one mechanism is necessary to achieve the same end. We show here that other uncoupling processes cannot completely compensate for a loss of MSL1: the mitochondrial glutathione pool (as indicated by the degree of mito-roGFP2 oxidation) is more oxidised in the *msl1-1* mutant under specific

abiotic stress conditions and when the mitochondrial bioenergetic state is directly manipulated with respiratory inhibitors (Figure 7b). This is despite the observation that loss of MSL1 causes an increase in pulsing (Figure 5). This suggests that MSL1, pulsing and UCP are non-redundant *in planta*.

Three reasons can be envisaged for multiple routes to mitochondrial uncoupling. Firstly, it is possible that different mechanisms could lead to activation at different levels of mitochondrial respiratory chain over-reduction. The second reason could be related to capacity in different cell types. The abundance of the transporters and channels responsible could be quite different leading to different capacities for uncoupling. Combined with different activation thresholds, this would allow a fine tuning of the response, which may be important to dampen bioenergetic imbalance while minimising the effect on ATP synthesis. Finally, it is evident that the different uncoupling mechanisms operate on different timescales, and this could be important in allowing both short- and long-term homeostasis and adaptation to environment. In particular, the alternative respiratory pathways appear to be largely regulated at the transcriptional level (Escobar *et al.*, 2004; Clifton *et al.*, 2005), which means they allow relatively long-term adjustments to environmental stress. In contrast, pulsing operates on the timescale of less than 1 min (Schwarzländer *et al.*, 2012), and MSL1 function is also likely to be rapid given the timescale of its response to bioenergetic state changes in isolated mitochondria. It is possible that MSL1 may function to limit increases in mitochondrial membrane potential prior to the onset of pulsing.

In summary, we have shown that MSL1 is a stretch-activated ion channel with electrophysiological similarities to its homologues from *E. coli* and Arabidopsis. While mitochondrially localised MSL1 can confer protection from osmotic shock onto *E. coli* lacking key bacterial mechanosensitive channels, it does not protect isolated mitochondria against lysis under hypo-osmotic shock. Instead, we demonstrate that MSL1 function is apparent when mitochondrial membrane potential is high, which is also a condition that leads to mitochondrial ROS production and lipid peroxidation. We also show that the absence of MSL1 loads a greater oxidative burden on mitochondria under abiotic stress, reflected in an increased oxidation of the mitochondrial glutathione pool.

EXPERIMENTAL PROCEDURES

Plant materials and growth conditions

Arabidopsis thaliana (ecotype Columbia-0) seedlings and plants were grown under a light intensity of $50 \mu\text{mol m}^{-2} \text{sec}^{-1}$ with a photoperiod of 16 h after seeds were stratified at 4°C in the dark for 2–3 days. The *msl1-1* T-DNA insertion line was obtained directly from GABI-Kat (GK-844G07). Plants were grown on

compost supplemented with Vermiculite. For the isolation of mitochondria, seeds were surface-sterilised and grown in Murashige and Skoog (MS) medium (½-strength MS medium, 2 mM MES pH 5.7) supplemented with 1% (w/v) sucrose and 0.1% (w/v) agar for 14–16 days with gentle agitation (40–60 rpm). For growth assays or *in vivo* CLSM of roots, seedlings were grown for 5–6 days on vertical plates containing ½-MS medium and 1.5% (w/v) agar.

Generation of MSL1 transgenic lines and mutant identification

Plants homozygous for a T-DNA insertion in the first exon of *MSL1* were identified by PCR of genomic DNA using the following primer pairs: MSL1_F and MSL1_R; T-DNA_F and T-DNA_R (Table S1). To visually confirm the absence of *MSL1* transcript in the insertion line, cDNA was synthesised from DNase-treated total RNA as previously described (Birke *et al.*, 2013), and semi-quantitative PCR was carried out with the primer pair MSL1_sqPCR_F and MSL1_sqPCR_R (Table S1). Quantitative RT-PCR was performed in a Rotor-Gene Q cyclor (Qiagen, Düsseldorf, Germany), using the Rotor-Gene SYBR Green PCR kit (Qiagen) according to the manufacturer's instructions with the primers MSL1_qPCR_F and MSL1_qPCR_R (Table S1). The amount of MSL1 protein in the transgenic lines was quantified by immunoblotting using a peptide antibody raised against the C-terminus of the MSL1 protein (see below).

The *MSL1* genomic complementation transgene *MSL1 g* was made by amplifying the entire *MSL1* genomic region, including the promoter, 5', and 3' UTRs from WT Columbia DNA with oligos LHO710 and LHO817 (Table S1). The resulting 3.6-kb fragment was cloned via the Gateway system into pENTR/D-TOPO and then into the destination binary vector pBGW. The complemented lines were generated by Agrobacterium-mediated transformation of the pBGW-MSL1g construct. Primary transformants were screened for BASTA resistance and the presence of *MSL1* transcript by RT-PCR. After two generations (T_2), a segregation analysis was performed and the homozygous complemented line with an approximately equal amount of *MSL1* transcript as Col-0 was chosen for further analysis. For generating the MSL1-GFP construct, the *MSL1* open reading frame was amplified from leaf cDNA with MSL1-GFP_F and MSL1-GFP_R (Table S1), and cloned via the Gateway system into pENTR/D-TOPO and then into the destination binary vector pK7FWG2. This construct was then introduced into Col-0 by Agrobacterium-mediated transformation. To generate the *msl1-1/roGFP2* line, the homozygous *msl1-1* line was crossed with a transgenic line carrying a roGFP2 construct (Schwarzländer *et al.*, 2008).

Isolation and fractionation of mitochondria

Mitochondria were isolated from 2-week-old Arabidopsis seedlings as described previously (Sweetlove *et al.*, 2007). Fractionation of mitochondria into peripheral and integral membrane fractions was carried out according to Millar *et al.* (2001) with slight modifications. Briefly, 1 mg of mitochondrial proteins was resuspended in 100 μl of 20 mM TES (2-[(2-hydroxy-1,1-bis(hydroxymethyl)ethyl)amino]ethanesulfonic acid) buffer (pH 7.5) and incubated for 10 min. Following three freeze-thaw cycles, the pellet containing membrane-associated proteins was collected by centrifugation at 20 000 *g* for 25 min. It was then resuspended in 100 μl of 100 mM Na_2CO_3 (pH 11.5) and incubated on ice for 20 min. The resulting sample was centrifuged at 20 000 *g* to collect the peripheral and integral membrane fractions in the

supernatant and pellet, respectively. The pellet was resuspended in blue-native solubilisation buffer containing digitonin before loading onto a blue native- or SDS-PAGE.

Mitoplasts were obtained by hypo-osmotic shock of mitochondria, using a method described previously (Murcha *et al.*, 2003). To determine the topology of proteins, 750 µg of mitochondria or mitoplasts were treated with 24 µg ml⁻¹ proteinase K (Sigma-Aldrich, Gillingham, UK) or water on ice. Following 30 min incubation, 1 mM PMSF was added to the samples to stop the reaction. Mitochondria or mitoplasts were collected by sedimentation at 20 000 g for 20 min. This step effectively removed un-pelleted proteins in the intermembrane space in the mitoplast fraction. The pellet was resuspended in 75 µl of 100 mM Na₂CO₃ (pH 11.5), and integral membrane proteins were extracted as described above. *In vitro* import of MSL1 into isolated mitochondria and chloroplasts was carried as described previously (Duncan *et al.*, 2015).

CLSM and data analysis

Freshly isolated mitochondria (in 1 mg ml⁻¹) were incubated in 500 nM MitoTracker Green in minimum medium [0.3 M sucrose, 10 mM TES and pH 7.2 (adjusted with NaOH)] for 30 min at 4°C. Mitochondria were then immobilised onto round glass cover slides by centrifugation in minimum medium at 2000 g for 5 min at 4°C with brake off. Immobilised mitochondria were energised by the addition of a minimum medium containing respiratory substrates and/or effectors (as stated in the text and in the figure legends) and labeled with 25 nM TMRM for state II respiration or 100 nM TMRM for state III respiration. Microscopy was performed using a Zeiss LSM510 META confocal microscope (Carl Zeiss Microimaging, Cambridge, UK) equipped with laser lines for 488- and 543-nm excitation. Images were collected with a 60× lens (Zeiss 60× 1.4 N.A. Plan-Apochromat oil-immersion lens) for multi-track mode with line switching between channels. Excitation/emission wavelengths were selected for different probes as follows: TMRM, 543 nm/565–615 nm; and MitoTracker Green, 488 nm/505–530 nm. The pinhole diameter was 1.2–1.4 airy units depending on the emission band-pass filter, but the optical slice was kept constant at 1.0 µm for all channels. All other imaging parameters, including laser power of both excitation channels, were kept constant throughout all measurements. CLSM z-stack (10 × 0.2 µm) and time series data (3 sec interval in the total of 2–3 min) of mitochondria were analyzed using a custom Matlab program available from www.markfricker.org (Fricker, 2016). In brief, the mitochondria were automatically separated into unique domains using a watershed algorithm and individually segmented with a local intensity threshold set at the mid-point between each object and the background. This was required to ensure that objects with different intensity values were segmented correctly, which was not possible with a single, global intensity threshold. The average fluorescence intensity at each wavelength was measured for each mitochondrion, and the volume estimated from the projected area assuming spherical geometry.

Confocal laser-scanning microscope imaging of MSL1-GFP seedlings was performed using a FLUOVIEW FV1000 (Olympus, Center Valley, PA, USA), and images were captured with FVIO-ASW software (Olympus). GFP signal was excited at 488 nm and emissions collected with a 505–520 nm bandpass filter. MitoTracker was excited at 543 nm and emissions collected with 560–620 nm BP filter. Chlorophyll autofluorescence was excited at 635 nm and emissions collected with a 655–755 nm BP filter.

For ratiometric *in vivo* imaging of 5–6-day-old roGFP2-expressing seedlings, a Nikon A1Si confocal microscope equipped with laser lines 405- and 488-nm excitation and emission band-pass

filter of 500–520 nm, and controlled by a NIS-Elements AR software package (version 4.13.01, Build 916) was used. Images were acquired in the '2Ex 1Em' mode using a 20× lens (Nikon CFI Plan Apo VC 20× 0.75 N.A.) with pinhole diameter of 2.5 airy units (corresponds to the optical slice of 4.37 µm). The 405/488-nm laser power ratio of 6:1 for imaging roGFP2 was kept constant for all treatments as indicated. The 405/488-nm fluorescence ratios and the degree of oxidation of roGFP2 were calculated using the same custom Matlab program described above (available from www.markfricker.org; Fricker, 2016). In brief, this approach finds the brightest voxel in z with excitation at 488 nm, and then collects the equivalent voxel from the 405 and autofluorescence channels before background subtraction and ratioing (Fricker, 2016). Calibration of the roGFP2 probe was carried out using 10 mM DTT (full reduction) and 10 mM H₂O₂ (full oxidation), and the degree of oxidation of roGFP2 was calculated according to Schwarzländer *et al.* (2008).

Molecular cloning and expression of MSL1

The gene corresponding to the mature version of MSL1 (i.e. without the first 79 amino acid residues that constitute the mitochondrial targeting sequence) was amplified with the primer pair His-MSL1-m_F and His-MSL1-m_R (Table S1). The PCR product was then subcloned into an entry vector using a pENTR/D-TOPO cloning kit. MSL1 was then cloned into a Champion pET300/NT-DEST vector (Life Technologies, Paisley, UK) by Gateway LR reaction, and the resulting construct was transformed into *E. coli* Rosetta strain. Before induction, bacteria were inoculated in 40 ml LB medium containing 1% glucose, 200 µg ml⁻¹ ampicillin and 34 µg ml⁻¹ chlorophenicol. Following overnight pre-induction, cells were collected by centrifugation (4000 g for 15 min), and the resulting pellet was resuspended in fresh LB-medium. Bacteria were added to 200 ml LB-medium containing 1% glucose and antibiotics to achieve the final OD_{550 nm} of at least 0.4. Induction was carried out overnight at 28°C after the addition of 1 mM IPTG. Cells were harvested by centrifugation (4000 g for 20 min), and the pellet was resuspended and clarified in 10 ml lysis buffer (50 mM phosphate buffer pH 8.0, 300 mM NaCl and 10 mM imidazole) containing 0.5% (v/v) Triton X-100. Following centrifugation to remove insoluble materials (e.g. inclusion bodies), proteins in the supernatant were separated overnight by a 4.5–16% BN-PAGE, and the identity of the MSL1 band was confirmed by immunoblotting and mass spectrometry (see below).

Inserts for pCTC-His6-MSL1 and pCTC-MSL1-FLAG were amplified from pET300-MSL1 using LHO2458 and LHO2459 or LHO2460 and LHO2462, respectively (Table S1). PCR products were inserted into pCTC-FLAG (Sigma) digested with EcoRI and Sall using a Gibson Assembly Cloning Kit (NEB).

Polyacrylamide gel electrophoresis, immunoblotting and mass spectrometry

To produce a specific antibody against AtMSL1, a homology modeling was performed, using the known structure of *E. coli* MscS (Bass *et al.*, 2002; Steinbacher *et al.*, 2007) as a template, to determine the peptide sequence where it is exposed to the aqueous environment that allows recognition and binding by the epitopes. Antibodies against the peptide sequence QISNEIKEMLRNNTK were produced in rabbits (Eurogentec, Seraing, Belgium). Purified antibodies were used in 1:1000 for probing the native form of MSL1.

Separation of digitonin-solubilised mitochondrial proteins was carried out according to Eubel *et al.* (2003) using a 4.5–16%

gradient BN gel dimension and a two-step Tricine-SDS second dimension (8 and 12%). For Western blotting, BN gels and PVDF membranes were equilibrated in a modified transfer buffer (48 mM Tris, 15 mM bis-Tris and 2.5 mM Tricine at 4°C). Protein transfer was carried out using a semi-dry FastBlot B43 system (Biometra, Göttingen, Germany) at 0.5 mA per cm² of gel for 3 h at 4°C.

For Western blotting with 1D SDS-PAGE, 20 µg proteins were separated by SDS-PAGE using a Hoefer MiniEV systems, and transferred to PVDF or nitrocellulose membranes for immunodetection by standard procedures using horseradish peroxidase-conjugated secondary antibody. Transferred proteins were probed with primary antibodies specifically targeted to MSL1 (1:1000), Porin (1:5000; Lee *et al.*, 2008), SLP2 (1:5000; Gehl *et al.*, 2014), cytochrome c (1:1000; Balk *et al.*, 1999) and penta-His (Anti-His Antibody Selector Kit, Qiagen).

For protein identification, gel pieces were destained in 25 mM ammonium bicarbonate in 50% acetonitrile, reduced (10 mM DTT, 37°C, 30 min) and alkylated (55 mM iodoacetamide, 25°C, 1 h). Trypsin digest [200 ng (Promega, Southampton, UK) in 25 mM ammonium bicarbonate] was performed at 37°C overnight. Extracted peptides were concentrated in a Savant SpeedVac (Thermo Fisher Scientific, Renfrew, UK). Samples were then processed for protein identification at the Central Proteomics Facility, Dunn School of Pathology, University of Oxford. Briefly, samples were resolved on a 25 cm × 75 µm diameter column in an Ultimate 3000 nanoRSLC system (Dionex, Hemel Hempstead, UK) at 300 nl min⁻¹ flow rate using a 125-min gradient before detection in a Q Exactive mass spectrometer (Thermo Fisher Scientific). Sample data were queried using Mascot Search Engine (version 2.4, MatrixScience, London, UK) against UniProtCP_arath. Search criteria were trypsin with two missed cleavage sites allowed; fixed modification was carbamidomethyl (C), variable modification was oxidised (M). Precursor mass tolerance was ±20 ppm and fragment mass tolerance was ±0.02 Da.

Osmotic shock assays

Bacterial osmotic downshock assays were performed essentially as described in Bartlett *et al.* (2004). Cultures of bacteria (transformed either with a pCTC-FLAG, pCTC-His6-MSL1 or pCTC-MSL1-FLAG construct) were inoculated from newly transformed colonies and grown overnight at 37°C, then diluted 1:10 and grown to an OD₆₀₀ of 0.1. IPTG induction was for 2 h.

Electrophysiology

Giant spheroplasts were made as described in Martinac *et al.* (2013), with the following modifications. *Escherichia coli* cell cultures were treated with cephalixin for 1.5 h and then induced with 1 mM IPTG for 2 h. Spheroplasting was carried out at room temperature for 20 min. The resulting suspension was centrifuged through 7 ml of 1 M sucrose, resuspended in 1 M sucrose, and stored in aliquots at -20°C or -80°C.

Patch-clamp experiments were performed with pipette buffer (200 mM KCl, 90 mM MgCl₂, 5 mM CaCl₂, 5 mM Hepes, pH 7.4) and a bath buffer (pipette with 400 mM sucrose). Excised inside-out patches [generated with pipettes with bubble number ~4.5 (Schnorf *et al.* (1994))] from spheroplast membranes were treated with 5-sec triangle pressure ramps of amplitudes from -40 to -200 mmHg. Data were collected using an Axopatch 200B amplifier and a Digidata 1440 digitiser (Molecular Devices, Sunnyvale, CA, USA) at 20 kHz and filtered at 5 kHz. Data were analyzed with Clampfit 10.5 (pCLAMP 10 software suite, Molecular Devices).

The ion selectivity of MSL1 was probed under asymmetric buffer conditions. Inside-out patches were excised in symmetric buffer (200 mM KCl, 90 mM MgCl₂, 5 mM CaCl₂, 5 mM Hepes, pH 7.3). Then the bath was perfused with a high salt solution (600 mM KCl, 90 mM MgCl₂, 5 mM CaCl₂, 5 mM Hepes, pH 7.3). The value of reversal potential was calculated as the point of intersection of the voltage axis with a line representing a linear fit of unitary current in the range between 40 mV and -50 mV membrane potential.

ACKNOWLEDGEMENTS

CPL was supported by a European Molecular Biology Organization (EMBO) long-term fellowship (ALTF 1140-2011). CPL also acknowledges support from the CellNetworks Excellence Cluster Postdoctoral Program (University of Heidelberg) to initiate this work. AHM was supported by the ARC (Australian Research Council) Centre of Excellence in Plant Energy Biology (grant numbers CE140100008; FT110100242). ESH acknowledges the support of the NIH (National Institutes of Health) grant number 2R01GM084211-5A1. The authors also acknowledge the Centre for Microscopy, Characterisation & Analysis (CMCA) at the University of Western Australia for the use of facilities and technical assistance. Finally, the authors would like to thank Mike Murphy (Medical Research Council Mitochondrial Unit, Cambridge, UK) for providing MitoQ.

SUPPORTING INFORMATION

Additional Supporting Information may be found in the online version of this article.

Figure S1. The MSL1 precursor is not imported into isolated chloroplasts *in vitro*.

Figure S2. Characterisation of *msl1-1*-knockout and complemented lines of Arabidopsis.

Figure S3. Verification of MSL1 expression in *Escherichia coli*.

Figure S4. Comparison of tension-gated MSL1 channel activity under positive and negative membrane potentials.

Data S1. Identification of MSL1 by mass spectrometry.

Table S1. A list of primers used in this study.

REFERENCES

- Atkin, O.K., Zhang, Q. and Wiskich, J.T. (2002) Effect of temperature on rates of alternative and cytochrome pathway respiration and their relationship with the redox poise of the quinone pool. *Plant Physiol.* **128**, 212–222.
- Azzu, V., Jastroch, M., Divakaruni, A.S. and Brand, M.D. (2010) The regulation and turnover of mitochondrial uncoupling proteins. *Biochim. Biophys. Acta*, **1797**, 785–791.
- Balk, J., Leaver, C.J. and McCabe, P.F. (1999) Translocation of cytochrome c from the mitochondria to the cytosol occurs during heat-induced programmed cell death in cucumber plants. *FEBS Lett.* **463**, 151–154.
- Bar-On, P., Mohsen, M., Zhang, R., Feigin, E., Chevion, M. and Samuni, A. (1999) Kinetics of nitroxide reaction with iron(II). *J. Am. Chem. Soc.* **121**, 8070–8073.
- Barreto, P., Okura, V., Neshich, I.A., Maia, I.D. and Arruda, P. (2014) Overexpression of UCP1 in tobacco induces mitochondrial biogenesis and amplifies a broad stress response. *BMC Plant Biol.* **14**, 144.
- Bartlett, J.L., Levin, G. and Blount, P. (2004) An *in vivo* assay identifies changes in residue accessibility on mechanosensitive channel gating. *Proc. Natl Acad. Sci. USA*, **101**, 10161–10165.
- Bass, R.B., Strop, P., Barclay, M. and Rees, D.C. (2002) Crystal structure of *Escherichia coli* MscS, a voltage-modulated and mechanosensitive channel. *Science*, **298**, 1582–1587.
- Beavis, A.D. (1992) Properties of the inner membrane anion channel in intact mitochondria. *J. Bioenerg. Biomembr.* **24**, 77–90.

- Birke, H., Heeg, C., Wirtz, M. and Hell, R. (2013) Successful fertilization requires the presence of at least one major *O*-acetylserine(thio)lyase for cysteine synthesis in pollen of *Arabidopsis*. *Plant Physiol.* **163**, 959–972.
- Booth, I.R., Rasmussen, T., Edwards, M.D., Black, S., Rasmussen, A., Bartlett, W. and Miller, S. (2011) Sensing bilayer tension: bacterial mechanosensitive channels and their gating mechanisms. *Biochem. Soc. Trans.* **39**, 733–740.
- Clifton, R., Lister, R., Parker, K., Sappl, P., Elhafez, D., Millar, A.H., Day, D. and Whelan, J. (2005) Stress-induced co-expression of alternative respiratory chain components in *Arabidopsis thaliana*. *Plant Mol. Biol.* **58**, 193–212.
- Cox, C.D., Nomura, T., Ziegler, C.S., Campbell, A.K., Wann, K.T. and Martiniac, B. (2013) Selectivity mechanism of the mechanosensitive channel MscS revealed by probing channel subconducting states. *Nat. Commun.* **4**, 2137.
- Dioloz, P. and Moreau, F. (1985) Correlation between ATP synthesis, membrane potential and oxidation rate in potato mitochondria. *Biochim. Biophys. Acta*, **806**, 56–63.
- Duncan, O., Carrie, C., Wang, Y. and Murcha, M. (2015) *In vitro* and *in vivo* protein uptake studies in plant mitochondria. In *Plant Mitochondria* (Whelan, J. and Murcha, M.W., eds). New York: Springer, pp. 61–81.
- Edwards, M.D., Bartlett, W. and Booth, I.R. (2008) Pore mutations of the *Escherichia coli* MscS channel affect desensitization but not ionic preference. *Biophys. J.* **94**, 3003–3013.
- Edwards, M.D., Black, S., Rasmussen, T., Rasmussen, A., Stokes, N.R., Stephen, T.-L., Miller, S. and Booth, I.R. (2012) Characterization of three novel mechanosensitive channel activities in *Escherichia coli*. *Channels*, **6**, 272–281.
- Escobar, M.A., Franklin, K.A., Svensson, Å.S., Salter, M.G., Whitlam, G.C. and Rasmussen, A.G. (2004) Light regulation of the *Arabidopsis* respiratory chain. Multiple discrete photoreceptor responses contribute to induction of type II NAD(P)H dehydrogenase genes. *Plant Physiol.* **136**, 2710–2721.
- Eubel, H., Jansch, L. and Braun, H.-P. (2003) New insights into the respiratory chain of plant mitochondria. Supercomplexes and a unique composition of Complex II. *Plant Physiol.* **133**, 274–286.
- Ferro, M., Brugière, S., Salvi, D. et al. (2010) AT_CHLORO, a comprehensive chloroplast proteome database with subplastidial localization and curated information on envelope proteins. *Mol. Cell Proteomics*, **9**, 1063–1084.
- Fricke, M.D. (2016) Quantitative redox imaging software. *Antioxid. Redox Signal.* **24**, 752–762.
- Froehlich, J.E., Wilkerson, C.G., Ray, W.K., McAndrew, R.S., Osteryoung, K.W., Gage, D.A. and Phinney, B.S. (2003) Proteomic study of the *Arabidopsis thaliana* chloroplast envelope membrane utilizing alternatives to traditional two-dimensional electrophoresis. *J. Proteome Res.* **2**, 413–425.
- Gehl, B., Lee, C.P., Bota, P., Blatt, M.R. and Sweetlove, L.J. (2014) An *Arabidopsis* stomatin-like protein affects mitochondrial respiratory supercomplex organization. *Plant Physiol.* **164**, 1389–1400.
- Hamilton, E.S., Schlegel, A.M. and Haswell, E.S. (2015a) United in diversity: mechanosensitive ion channels in plants. *Annu. Rev. Plant Biol.* **66**, 113–137.
- Hamilton, E.S., Jensen, G.S., Maskaev, G., Katims, A., Sherp, A.M. and Haswell, E.S. (2015b) Mechanosensitive channel MSL8 regulates osmotic forces during pollen hydration and germination. *Science*, **350**, 438–441.
- Haswell, E.S. and Meyerowitz, E.M. (2006) MscS-like proteins control plastid size and shape in *Arabidopsis thaliana*. *Curr. Biol.* **16**, 1–11.
- Hensley, J.R. and Hanson, J.B. (1975) The action of valinomycin in uncoupling corn mitochondria. *Plant Physiol.* **56**, 13–18.
- Huang, S., Taylor, N.L., Whelan, J. and Millar, A.H. (2009) Refining the definition of plant mitochondrial presequences through analysis of sorting signals, N-terminal modifications, and cleavage motifs. *Plant Physiol.* **150**, 1272–1285.
- Huber, S.C. and Moreland, D.E. (1979) Permeability properties of the inner membrane of mung bean mitochondria and changes during energization. *Plant Physiol.* **64**, 115–119.
- Jezek, P., Costa, A.D.T. and Vercesi, A.E. (1997) Reconstituted plant uncoupling mitochondrial protein allows for proton translocation via fatty acid cycling mechanism. *J. Biol. Chem.* **272**, 24272–24278.
- Jung, D.W. and Brierley, G.P. (1979) Swelling and contraction of potato mitochondria. *Plant Physiol.* **64**, 948–953.
- Kelso, G.F., Porteous, C.M., Coulter, C.V., Hughes, G., Porteous, W.K., Ledgerwood, E.C., Smith, R.A.J. and Murphy, M.P. (2001) Selective targeting of a redox-active ubiquinone to mitochondria within cells: antioxidant and antiapoptotic properties. *J. Biol. Chem.* **276**, 4588–4596.
- Kessler, A. and Brand, M.D. (1994a) Effects of cadmium on the control and internal regulation of oxidative phosphorylation in potato tuber mitochondria. *Eur. J. Biochem.* **225**, 907–922.
- Kessler, A. and Brand, M.D. (1994b) Quantitative determination of the regulation of oxidative phosphorylation by cadmium in potato tuber mitochondria. *Eur. J. Biochem.* **225**, 923–935.
- Kessler, A. and Brand, M.D. (1994c) Localisation of the sites of action of cadmium on oxidative phosphorylation in potato tuber mitochondria using top-down elasticity analysis. *Eur. J. Biochem.* **225**, 897–906.
- Klodmann, J., Senkler, M., Rode, C. and Braun, H.-P. (2011) Defining the protein complex proteome of plant mitochondria. *Plant Physiol.* **157**, 587–598.
- Kung, C., Martinac, B. and Sukharev, S. (2010) Mechanosensitive channels in microbes. *Annu. Rev. Microbiol.* **64**, 313–329.
- Kunz, H.H., Gierth, M., Herdean, A., Satoh-Cruz, M., Kramer, D.M., Spetea, C. and Schroeder, J.I. (2014) Plastidial transporters KEA1, -2, and -3 are essential for chloroplast osmoregulation, integrity, and pH regulation in *Arabidopsis*. *Proc. Natl Acad. Sci. USA*, **111**, 7480–7485.
- Laus, M.N., Soccio, M., Trono, D., Cattivelli, L. and Pastore, D. (2008) Plant inner membrane anion channel (PIMAC) function in plant mitochondria. *Plant Cell Physiol.* **49**, 1039–1055.
- Lee, C.P., Eubel, H., O'Toole, N. and Millar, A.H. (2008) Heterogeneity of the mitochondrial proteome for photosynthetic and non-photosynthetic *Arabidopsis* metabolism. *Mol. Cell Proteomics*, **7**, 1297–1316.
- Levina, N., Töttemeyer, S., Stokes, N.R., Louis, P., Jones, M.A. and Booth, I.R. (1999) Protection of *Escherichia coli* cells against extreme turgor by activation of MscS and MscL mechanosensitive channels: identification of genes required for MscS activity. *EMBO J.* **18**, 1730–1737.
- Li, Y., Moe, P.C., Chandrasekaran, S., Booth, I.R. and Blount, P. (2002) Ionic regulation of MscK, a mechanosensitive channel from *Escherichia coli*. *EMBO J.* **21**, 5323–5330.
- Lin, T.-Y. and Markhart, A.H. (1990) Temperature effects on mitochondrial respiration in *Phaseolus acutifolius* A. Gray and *Phaseolus vulgaris* L. *Plant Physiol.* **94**, 54–58.
- Maskaev, G. and Haswell, E.S. (2012) MscS-Like10 is a stretch-activated ion channel from *Arabidopsis thaliana* with a preference for anions. *Proc. Natl Acad. Sci. USA*, **109**, 19015–19020.
- Martinac, B., Rohde, P., Cranfield, C. and Nomura, T. (2013) Patch clamp electrophysiology for the study of bacterial ion channels in giant spheroplasts of *E. coli*. In *Bacterial Cell Surfaces* (Delcour, A.H., ed.). New York City: Humana Press, pp. 367–380.
- Millar, A.H., Sweetlove, L.J., Giegé, P. and Leaver, C.J. (2001) Analysis of the *Arabidopsis* mitochondrial proteome. *Plant Physiol.* **127**, 1711–1727.
- Miller, S., Bartlett, W., Chandrasekaran, S., Simpson, S., Edwards, M. and Booth, I.R. (2003) Domain organization of the MscS mechanosensitive channel of *Escherichia coli*. *EMBO J.* **22**, 36–46.
- Mitchell, P. and Moyle, J. (1969) Estimation of membrane potential and pH difference across the cristae membrane of rat liver mitochondria. *Eur. J. Biochem.* **7**, 471–484.
- Møller, I.M. (2001) Plant mitochondria and oxidative stress: electron transport, nadph turnover, and metabolism of reactive oxygen species. *Annu. Rev. Plant Physiol. Plant Mol. Biol.* **52**, 561–591.
- Moore, A.L. and Wilson, S.B. (1977) Translocation of some anions, cations, and acids in turnip (*Brassica napus* L.) mitochondria. *J. Exp. Bot.* **28**, 607–618.
- Moore, A.L., Shiba, T., Young, L., Harada, S., Kita, K. and Ito, K. (2013) Unraveling the heater: new insights into the structure of the alternative oxidase. *Annu. Rev. Plant Biol.* **64**, 637–663.
- Murcha, M.W., Lister, R., Ho, A.Y.Y. and Whelan, J. (2003) Identification, expression, and import of components 17 and 23 of the inner mitochondrial membrane translocase from *Arabidopsis*. *Plant Physiol.* **131**, 1737–1747.
- Murcha, M.W., Elhafez, D., Millar, A.H. and Whelan, J. (2004) The N-terminal extension of plant mitochondrial carrier proteins is removed by two-step

- processing: the first cleavage is by the mitochondrial processing peptidase. *J. Mol. Biol.* **344**, 443–454.
- Murphy, M.P. and Smith, R.A.J.** (2007) Targeting antioxidants to mitochondria by conjugation to lipophilic cations. *Annu. Rev. Pharmacol. Toxicol.* **47**, 629–656.
- Naismith, J.H. and Booth, I.R.** (2012) Bacterial mechanosensitive channels – MscS: evolution's solution to creating sensitivity in function. *Annu. Rev. Biophys.* **41**, 157–177.
- Nicholls, D.G.** (1974) The influence of respiration and ATP hydrolysis on the proton-electrochemical gradient across the inner membrane of rat-liver mitochondria as determined by ion distribution. *Eur. J. Biochem.* **50**, 305–315.
- Nikolovski, N., Rubtsov, D., Segura, M.P., Miles, G.P., Stevens, T.J., Dunkley, T.P.J., Munro, S., Lilley, K.S. and Dupree, P.** (2012) Putative glycosyltransferases and other plant golgi apparatus proteins are revealed by LOPIT proteomics. *Plant Physiol.* **160**, 1037–1051.
- Pivetti, C.D., Yen, M.R., Miller, S., Busch, W., Tseng, Y.H., Booth, I.R. and Saier, M.H. Jr** (2003) Two families of mechanosensitive channel proteins. *Microbiol. Mol. Biol. Rev.* **67**, 66–85.
- Quinlan, C.L., Gerencser, A.A., Treberg, J.R. and Brand, M.D.** (2011) The mechanism of superoxide production by the antimycin-inhibited mitochondrial Q-cycle. *J. Biol. Chem.* **286**, 31361–31372.
- Rasmusson, A.G., Soole, K.L. and Elthon, T.E.** (2004) Alternative NAD(P)H dehydrogenases of plant mitochondria. *Annu. Rev. Plant Biol.* **55**, 23–39.
- Ridone, P., Nakayama, Y., Martinac, B. and Battle, A.R.** (2015) Patch clamp characterization of the effect of cardiolipin on MscS of *E. coli*. *Eur. Biophys. J.* **44**, 567–576.
- Samuni, A., Krishna, C.M., Mitchell, J.B., Collins, C.R. and Russo, A.** (1990) Superoxide reaction with nitroxides. *Free Radic Res. Commun.* **9**, 241–249.
- Santo-Domingo, J. and Demaurex, N.** (2012) The renaissance of mitochondrial pH. *J. Gen. Physiol.* **139**, 415–423.
- Schnorf, M., Potrykus, I. and Neuhaus, G.** (1994) Microinjection technique: routine system for characterization of microcapillaries by bubble pressure measurement. *Exp. Cell Res.* **210**, 260–267.
- Schwacke, R., Schneider, A., van der Graaff, E., Fischer, K., Catoni, E., Desimone, M., Frommer, W.B., Flügge, U.-I. and Kunze, R.** (2003) ARAMEMNON, a novel database for arabidopsis integral membrane proteins. *Plant Physiol.* **131**, 16–26.
- Schwarzländer, M., Fricker, M.D., Müller, C., Marty, L., Brach, T., Novak, J., Sweetlove, L.J., Hell, R. and Meyer, A.J.** (2008) Confocal imaging of glutathione redox potential in living plant cells. *J. Microsc.* **231**, 299–316.
- Schwarzländer, M., Logan, D.C., Johnston, I.G., Jones, N.S., Meyer, A.J., Fricker, M.D. and Sweetlove, L.J.** (2012) Pulsing of membrane potential in individual mitochondria: a stress-induced mechanism to regulate respiratory bioenergetics in Arabidopsis. *Plant Cell*, **24**, 1188–1201.
- Schwarzländer, M., Dick, T.P., Meyer, A.J. and Morgan, B.** (2016) Dissecting redox biology using fluorescent protein sensors. *Antioxid. Redox Signal.* **24**, 680–712.
- Smith, A.M.O., Ratcliffe, R.G. and Sweetlove, L.J.** (2004) Activation and function of mitochondrial uncoupling protein in plants. *J. Biol. Chem.* **279**, 51944–51952.
- Steinbacher, S., Bass, R.B., Strop, P. and Rees, D.C.** (2007) Structures of the prokaryotic mechanosensitive channels MscL and MscS. In *Mechanosensitive Ion Channels, Part A* (Hamill, O.P., ed.). Cambridge: Academic Press, pp. 1–24.
- Sukharev, S.** (2002) Purification of the small mechanosensitive channel of *Escherichia coli* (MscS): the subunit structure, conduction, and gating characteristics in liposomes. *Biophys. J.* **83**, 290–298.
- Sweetlove, L.J., Taylor, N.L. and Leaver, C.J.** (2007) Isolation of intact, functional mitochondria from the model plant *Arabidopsis thaliana*. *Methods Mol. Biol.* **372**, 125–136.
- Trnka, J., Blaikie, F.H., Logan, A., Smith, R.A.J. and Murphy, M.P.** (2009) Antioxidant properties of MitoTEMPOL and its hydroxylamine. *Free Radic. Res.* **43**, 4–12.
- Trono, D., Laus, M.N., Soccio, M., Alfarano, M. and Pastore, D.** (2015) Modulation of potassium channel activity in the balance of ROS and ATP production by durum wheat mitochondria – an amazing defense tool against hyperosmotic stress. *Front. Plant Sci.* **6**, 1072.
- Veley, K.M., Marshburn, S., Clure, C.E. and Haswell, E.S.** (2012) Mechanosensitive channels protect plastids from hypoosmotic stress during normal plant growth. *Curr. Biol.* **22**, 408–413.
- Veley, K.M., Maskaev, G., Frick, E.M., January, E., Kloepper, S.C. and Haswell, E.S.** (2014) Arabidopsis MSL10 has a regulated cell death signaling activity that is separable from its mechanosensitive ion channel activity. *Plant Cell*, **26**, 3115–3131.
- Vercesi, A.E., Borecký, J., Maia Ide, G., Arruda, P., Cuccovia, I.M. and Chaimovich, H.** (2006) Plant uncoupling mitochondrial proteins. *Annu. Rev. Plant Biol.* **57**, 383–404.
- Wagner, S., De Bortoli, S., Schwarzländer, M. and Szabó, I.** (2016) Regulation of mitochondrial calcium in plants versus animals. *J. Exp. Bot.*, **67**, 3809–3829.
- Wagner, S., Behera, S., De Bortoli, S. et al.** (2015) The EF-Hand Ca²⁺ binding protein MICU choreographs mitochondrial Ca²⁺ dynamics in Arabidopsis. *Plant Cell*, **27**, 3190–3212.
- Wilson, M.E., Jensen, G.S. and Haswell, E.S.** (2011) Two mechanosensitive channel homologs influence division ring placement in Arabidopsis chloroplasts. *Plant Cell*, **23**, 2939–2949.
- Wittig, I., Beckhaus, T., Wumaier, Z., Karas, M. and Schagger, H.** (2010) Mass estimation of native proteins by blue native electrophoresis: principles and practical hints. *Mol. Cell Proteomics*, **9**, 2149–2161.
- Zotova, L., Aleschko, M., Sponder, G., Baumgartner, R., Reipert, S., Prinz, M., Schweyen, R.J. and Nowikovsky, K.** (2010) Novel components of an active mitochondrial K⁺/H⁺ exchange. *J. Biol. Chem.* **285**, 14399–14414.
- Zybailov, B., Rutschow, H., Friso, G., Rudella, A., Emanuelsson, O., Sun, Q. and van Wijk, K.J.** (2008) Sorting signals, N-terminal modifications and abundance of the chloroplast proteome. *PLoS ONE*, **3**, e1994.